

Does Vitligo Affect the Gingiva?

Farah Khaled El-dernawi 1865 2nd year dentistry student

Libyan International Medical University



### Introduction

- ▶ Vitiligo or leukoderma is a skin disease with a world-▶ Very mild or no pigmentation in the epithelium wide prevalence ranging from 0.5% to 4%
- Characterized by partial or complete loss of pigment-producing melanocytes within the epidermis or mucosa
- Clinical lesions are asymptomatic, flat, well-demarcated macules and patches of pigment loss; their size varies from few to several centimeters, which may or may not be surrounded by macroscopic hy-.perpigmentation
  - Most noticeable in darkly pigmented individuals
- Vitiligo has been classified into three types, focal generalized, and segmental 1

### Histopathology

- ▶ Epidermal vacuolization
- Loose fibrous connective tissue with no evidence of inflammatory cells
- ► Melanocytes did not show immunoreactivity for HMB -45 antibody 4





normal gingival tissue

epithelium with vitiligo

## **Gingival Vitiligo**

- ▶ The incidence of vitiligo in the gingiva is rare
- It may occur only in the gingiva (Localized), or itmay be an effect of generalized vitiligo
- Vitiligo of the gingiva could result in cosmetic debilitywhich can in turn affect the mental status of the patient 4

# **2** Etiopathogenesis

#### i. Genetics

,Vitiligo is a typical polygenic multi factorial disorder, involving numerous different susceptibility genes

(HLA)(PTPN22)(CTLA4)(TYR) (TGFBR2)(UVRAG)(NLRP1) and

(SLEV1) on chromosome 17p13 a linkage signal in systemic lupus erythma tosus in generalized vitiligo 2



# CONCLUSION

### ii.Oxidative Stress

Vitiligo patients have an intrinsic defects that reduce the capacity of melanocytes to mitigate cellular stress ROS

#### iii.Environment

An exposure of phenolic and catecholic chemicals found in dyes, resins and leather may play a central role in disease onset

### iv.Immunity

Tyrosine hydroxylase(TH) antibodies(MCHR1) antibodies, IgM and IgG, anti-thyroglobulin antibodies and anti-thyroid antibodies. Cytotoxic CD8+ T cells are responsible for the destruction of melanocytes3

Vitiligo is a common pigmentary disorder, The melanocytes which produce melanin pigments become inactive as a result of genetic and environmental factors, oxidative stress, innate and adaptive immunity, Vitiligo can affect the gingiva and other oral mucosa

- .Your Bibliography: Forschner, T., Buchholtz, S., & Stockfleth, E 2007 Current state of vitiligo therapy? evidence-based analysis of the literature
- R Spritz 2011. The genetics of vitiligo
- Rashighi, M., & Harris, J 2017Vitiligo pathogenesis and emerging
- Ashok, N., Karunakaran, A., Singh, P., Rodrigues, J., Ashok, N .Tarakji, B. et al &2014 Gingival Vitiligo: Report of a case and review of the literature